


Stacking Cannonballs

Dr. Bob

Cannonballs in a layer

Let L_n be the number of cannonballs on layer n .

Each row in a layer has 1 more than the previous row:




$$L_n = 1 + 2 + 3 + 4 + \dots + n$$


$$L_n = \sum_{i=1}^n i$$

Cannonballs in a Layer

To find the sum of cannonballs in a layer graphically, rearrange the rows as like this:




Copy and rotate



n

Stack the two layers on top of each other




$n + 1$

The double layer has $2 * L_n = n * (n + 1)$, or

$$L_n = n * (n + 1) / 2 = \sum_{i=1}^n i$$

Stack the Cannonballs

Let T_N be the number of cannonballs in a stack of N layers.



$$T_N = 1 + 3 + 6 + 10 + \dots + L_N$$

$$T_N = \sum_{n=1}^N L_n$$

The answer to the original question:


When is the number above a layer = the number in the layer?

$$T_4 = (1 + 3 + 6) + 10 = 10 + 10$$


At layer 4.

Another way to count.

Suppose each layer was moved so that the point ball of each layer is right above the other.




Stack the layers.



From the side it looks like this.

Add up the cannonballs:



$$T_4 = 1 * 4 + 2 * 3 + 3 * 2 + 4 * 1 = \sum w * h$$

$$T_N = \sum_{n=1}^N n * (N + 1 - n) = (N + 1) \sum_{n=1}^N n - \sum_{n=1}^N n^2$$

Combine both ways.

For N layers, the last page gives:

$$T_N = \sum_{n=1}^N n * (N + 1 - n) = (N + 1) \sum_{n=1}^N n - \sum_{n=1}^N n^2 \quad (\text{eq. 1})$$

The first page gives:

$$T_N = \sum_{n=1}^N L_n = \sum_{n=1}^N \sum_{i=1}^n i = \sum_{n=1}^N n * (n + 1) / 2 = \sum_{n=1}^N n^2 / 2 + \sum_{n=1}^N n / 2$$

or

$$2 * T_N = \sum_{n=1}^N n + \sum_{n=1}^N n^2 \quad (\text{eq. 2})$$

Add eq. 2 + eq 1.

$$2 * T_N + T_N = 3 * T_N = (N + 2) \sum_{n=1}^N n = (N + 2) * N * (N + 1) / 2$$

or

$$T_N = N * \frac{(N + 1)}{2} * \frac{(N + 2)}{3}$$

Question:

When is the number above a layer = the number in the layer?

$$T_{N-1} = L_N$$

or

$$(N-1) * \frac{N}{2} * \frac{(N+1)}{3} = N * \frac{(N+1)}{2}$$

Dividing both sides by $N * ((N+1)/6)$ gives:

$$N-1 = 3$$

$$N = 4$$

Checking we have:

$$T_3 = 3 * (4/2) * (5/3) = 10$$

$$L_4 = 4 * (5/2) = 10$$

So the equations work!